arXiv:1206.2501v3 [math.PR] 6 May 2013

Noname manuscript No.
(will be inserted by the editor)

large deviation probabilities for sums of independent bounded random variables

Xiequan Fan · Ion Grama · Quansheng Liu

Received: date / Accepted: date

Abstract We obtain several optimal inequalities on tail probabilities for sums of independent and bounded random variables. The results complete an upper bound on tail probabilities due to Talagrand by giving a sharp lower bound and some explicit expressions for the constants of Talagrand's bound. In particular, we obtain an one-term asymptotic expansion for large deviations, which can be regarded as an inequality types of Cramér and Bahadur-Rao. The asymptotic expansion shows that the Talagrand inequality is sharp.

Keywords Sums of independent random variables · Talagrand's inequality · lower bound · Hoeffding's inequality · large deviations · random walk

Mathematics Subject Classification (2000) 60G50 · 60F10 · 60E15 · 62E20

1 Introduction

Let $\xi_1, ..., \xi_n$ be a sequence of random variables (r.v.s) satisfying $\mathbf{E}\xi_i = 0$. Denote $X_n = \sum_{i=1}^n \xi_i$. The study of tail probabilities $\mathbf{P}(X_n \geq x)$ has a long history. Interesting asymptotic expansions are obtained by Cramér (1938), Bahadur and Rao (1960), Petrov (1975, 1995), Saulis and Statulevičius (1991), Rozovky (2005), Bercu et al. (2006, 2011), Petrov and Robinson (2008) and Györfi, Harremöes and Tusnády (2012). Various upper bounds have been obtained: see Nagaev (1979), Pinelis and Utev (1989), Talagrand (1995, 1996a), Ledoux and Talagrand (1996) and Nagaev (2002) in the independent case; see Talagrand (1996b), Rio (2002, 2012) and van de Geer and Lederer (2011) for empirical processes; see Haeusler (1984), van de Geer (1995, 2002), Nagaev (2003, 2007), Bercu and Touati (2008) and Fan et al. (2012) for martingales; see Doukhan and Neumann (2007) and Delyon (2009) for sums of weakly dependent r.v.s.. A few results on lower bounds, we refer to Nagaev (2002) and Rozovky (2003) for sums of independent r.v.s.

In the paper, we consider the sums of independent and bounded r.v.s $(\xi_i)_{i=1,...,n}$. Denote by $\sigma_i^2 = \mathbf{E}\xi_i^2$ and $\sigma^2 = \sum_{i=1}^n \sigma_i^2$. Then it holds $\sigma = \sqrt{n}\,\sigma_1$ in the independent and identically distributed (i.i.d.) case. The celebrated Bennett inequality [2] states that: if $|\xi_i| \leq 1$ for all i,

---

X. Fan (✉) · I. Grama · Q. Liu
LMBA, CNRS UMR 6205, Université de Bretagne-Sud
E-mail: fanxiequan@hotmail.com; E-mail: ion.grama@univ-ubs.fr, quansheng.liu@univ-ubs.fr

then, for all $x > 0$,

$$\mathbf{P}(X_n \geq x\sigma) \leq B(x,\sigma) := \left(\frac{x+\sigma}{\sigma}\right)^{-\sigma x - \sigma^2} e^{x\sigma}. \tag{1}$$

However, Bennett's inequality (1) is not optimal. One of the most well-known improvements of (1) is Hoeffding's inequality (cf. (2.8) of [20]), which states that: if $\xi_i \leq 1$ for all i, then, for all $x > 0$,

$$\begin{aligned} \mathbf{P}(X_n \geq x\sigma) \ &\leq \ H_n(x,\sigma) && (2)\\ &:= \left\{ \left(\frac{\sigma}{x+\sigma}\right)^{x\sigma+\sigma^2} \left(\frac{n}{n-x\sigma}\right)^{n-x\sigma} \right\}^{\frac{n}{n+\sigma^2}} \mathbf{1}_{\{x \leq \frac{n}{\sigma}\}} && (3)\\ &\leq \ B(x,\sigma), && (4) \end{aligned}$$

where (and hereafter) by convention $(\frac{n}{n-x})^{n-x} = 1$ when $x = n$. By considering the following distribution

$$\mathbf{P}(\eta_i = 1) = \frac{\sigma^2/n}{1+\sigma^2/n} \quad \text{and} \quad \mathbf{P}(\eta_i = -\sigma^2/n) = \frac{1}{1+\sigma^2/n}, \tag{5}$$

Hoeffding showed that (2) is the best that can be obtained from the exponential Markov inequality

$$\mathbf{P}(X_n \geq x\sigma) \leq \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}, \qquad x \geq 0,$$

due to the element equality $\inf_{\lambda\geq 0} \mathbf{E}\exp\{\lambda(\sum_i^n \eta_i - x\sigma)\} = H_n(x,\sigma)$ for all $0 \leq x \leq \frac{n}{\sigma}$.

However, Hoeffding's inequality (2) can be still improved. For sums of bounded random variables, Talagrand [37] obtained the following improvement of (2). Assume that $(\xi_i)_{i=1,...,n}$ satisfy $-B \leq \xi_i \leq 1$ for some constant $B \geq 1$ and all i. Talagrand showed that, for all $0 \leq x \leq c_1 \frac{\sigma}{B}$,

$$\begin{aligned} \mathbf{P}(X_n \geq x\sigma) &\leq \left(\Theta(x) + c_2 \frac{B}{\sigma}\right) \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)} && (6)\\ &\leq \left(\Theta(x) + c_2 \frac{B}{\sigma}\right) H_n(x,\sigma), && (7) \end{aligned}$$

where $c_1, c_2 > 0$ are two absolute constants,

$$\Theta(x) = (1 - \Phi(x)) \exp\left\{\frac{x^2}{2}\right\}$$

is a decreasing function in $x \geq 0$ and $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{t^2}{2}} dt$ is the standard normal distribution function. Since

$$\frac{1}{\sqrt{2\pi}(1+x)} \leq \Theta(x) \leq \frac{1}{\sqrt{\pi}(1+x)}, \qquad x \geq 0, \tag{8}$$

(see Feller [15]), Talagrand's inequality (7) improves Hoeffding's inequality (2) by adding a factor of order $O(1/x)$ in the range $0 \leq x \leq c_1 \frac{\sigma}{B}$.

Since the inequalities of Talagrand are valid in the range $0 \leq x \leq c_1 \frac{\sigma}{B}$, it is important to know how large the constant c_1 can be to the practical use. For the sake of simplicity, consider the Rademacher. Denote by $S_n = \varepsilon_1 + ... + \varepsilon_n$ the sums of independent r.v.s ε_i with $\mathbf{P}(\varepsilon_i = \pm 1) = \frac{1}{2}$ for all $i = 1, ..., n$. Thus, we take $B = 1$ and $\sigma = \sqrt{n}$. The missing factor



Fig. 1 We display $\varTheta_n(x)$ and $\varTheta(x)$ as a function of x.

$$\varTheta_n(x) = \frac{\mathbf{P}(S_n \geq x\sqrt{n})}{H_n(x,\, \sqrt{n})} = \frac{\mathbf{P}(S_n \geq x\sqrt{n})}{\inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(S_n - x\sigma)}} \tag{9}$$

of Hoeffding's inequality (2) and $\varTheta(x)$ are displayed in Figure 1. From Figure 1, we find that the factor $\varTheta(x) + c_2\frac{B}{\sigma}$ in Talagrand's inequality (7) can dominate the missing factor $\varTheta_n(x)$ in Hoeffding's inequality for c_1 smaller than 1 (for instant $0 \leq x \leq 5$ in Figure 1). However, when c_1 is increasing to 1 (i.e. x increasing to 7), the constant c_2 in Talagrand's inequality (7) is increasing to a large number. In fact, since $\varTheta_n(\sqrt{n}) = 1$, from (7), we have $c_2 \to \sqrt{n}(1 - \varTheta(\sqrt{n})) \approx \sqrt{n}$ as $c_1 \to 1$. Thus, the constant c_2 in Talagrand's inequality (7) depends on c_1 and c_1 is less than 1.

In this paper, we give a sharp lower bound on tail probabilities and some information about c_1 and c_2 of Talagrand's inequality (6). Our main result Theorem 1 gives some explicit expressions for the constants c_1 and c_2 under a general assumption that $(\xi_i)_{i=1,\dots,n}$ have $(2+\delta)$th moments. In particular, Corollary 1 proves the following one-term asymptotic expansion (a type of one-term asymptotic expansion of Cramér [8] and Bahadur-Rao [1]): if $\xi_i \leq 1$ and $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ for some constant $B > 0$ and all i, then, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$,

$$\mathbf{P}(X_n \geq x\sigma) = \left[\varTheta(x) + 16\theta\frac{B}{\sigma}\right] \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}, \tag{10}$$

where $|\theta| \leq 1$. Since $\theta \geq -1$, we complete Talagrand's inequality (6) by giving a sharp lower bound. Taking into account the equality $\inf_{\lambda\geq 0} \mathbf{E}\exp\{\lambda(\sum_i^n \eta_i - x\sigma)\} = H_n(x, \sigma)$, we prove that Talagrand's inequality (7) is attained regardless of c_2. Moreover, by (8) and (10), we find that the related errors between $\mathbf{P}(X_n \geq x\sigma)$ and $\varTheta(x)\inf_{\lambda\geq 0}\mathbf{E}e^{\lambda(X_n - x\sigma)}$ converges to 0 uniformly in the range $0 \leq x = o(\frac{\sigma}{B})$ as $\frac{B}{\sigma} \to 0$,

$$\frac{\mathbf{P}(X_n \geq x\sigma)}{\varTheta(x)\inf_{\lambda\geq 0}\mathbf{E}e^{\lambda(X_n - x\sigma)}} = 1 + o(1). \tag{11}$$

To show the tightness of equality (11), we display the simulation of

$$R(x, n) = \frac{\mathbf{P}(S_n \geq x\sqrt{n})}{\varTheta(x)\inf_{\lambda\geq 0}\mathbf{E}e^{\lambda(S_n - x\sigma)}} = \frac{\mathbf{P}(S_n \geq x\sqrt{n})}{\varTheta(x)H_n(x, \sqrt{n})}$$



Fig. 2 The tail probabilities and ratios $R(x,n)$ are displayed as a function of x for various n.

in Figure 2, which shows that $R(x,n)$ is very close to 1 for large n's.

In the proofs of our paper, we make use of the technique based on the conjugate distribution technique, which become standard for obtaining Cramér large deviation results. Such results can be found in Cramér [8] and Petrov [31]. We refine the result of Talagrand [37] using a method inspired by Grama and Haeusler [16].

The paper is organized as follows. In Section 2, we present our main results. In Section 3, we present some auxiliary results. In Sections 4-6, we prove the main results.

Throughout the paper, write $a \wedge b = \min\{a,b\}$, $a \vee b = \max\{a,b\}$ and $a^+ = a \vee 0$. Denote by θ a number satisfying $|\theta| \leq 1$ and $\mathcal{N}(0,1)$ the standard normal random variable. We also agree that $0^{-1} = \infty$.

2 Main Results

Let $(\xi_i)_{i=1,\dots,n}$ be a sequence of independent real random variables with $\mathbf{E}\xi_i = 0$. In the sequel, we should use the following condition.

(A) There exist two constants $\delta \in (0,1]$ and $B > 0$ such that, for all $\lambda \geq 0$,

$$\mathbf{E}|\xi_i|^{2+\delta} \leq B^{2+\delta} \qquad \text{and} \qquad \sum_{i=1}^{n} \mathbf{E}\xi_i^2 e^{\lambda \xi_i} \geq (1 - B\lambda) \sum_{i=1}^{n} \mathbf{E}\xi_i^2 .$$

Notice that condition (A) is satisfied for $\delta = 1$ if $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ holds. Indeed, since $(\mathbf{E}\xi_i^2)^{3/2} \leq \mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$, we have $\mathbf{E}\xi_i^2 \leq B^2$ and $\mathbf{E}|\xi_i|^3 \leq B^3$. By the inequality $e^x \geq 1 + x$, it follows that $\sum_{i=1}^n \mathbf{E}\xi_i^2 e^{\lambda\xi_i} \geq \sum_{i=1}^n \mathbf{E}\xi_i^2(1+\lambda\xi_i) \geq (1-B\lambda)\sum_{i=1}^n \mathbf{E}\xi_i^2$ for all $\lambda \geq 0$.

For $\delta \in (0,1]$, denote by $C_{2+\delta}$ the Lyapunov constant defined as follows. Suppose that $(\xi_i)_{i=1,\dots,n}$ have $(2+\delta)$th moments, i.e. $\mathbf{E}|\xi_i|^{2+\delta} < \infty$ all i. Then $C_{2+\delta}$ is the minimum of all absolute constants C such that

$$\sup_{x\in\mathbf{R}} \Big| \mathbf{P}(X_n \leq x\sigma) - \Phi(x) \Big| \leq C\, \frac{\sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}}{\sigma^{2+\delta}}$$

holds for all $(\xi_i)_{i=1,\dots,n}$. It is known that $0.4097 \leq C_3 \leq 0.56$ and that $C_3 \leq 0.4784$ in the identically distributed case (see [36]). For the binomial distribution (for $0 < p \leq 0.5$), Nagaev and Chebotarev [26] recently have proved that $C_3 \leq 0.4215$.

Our first main result is the following theorem.

Theorem 1 *Assume that $(\xi_i)_{i=1,\dots,n}$ satisfy $\xi_i \leq 1$ and condition (A) for all i. Then, for all $0 \leq x < 0.25\frac{\sigma}{B}$,*

$$\mathbf{P}(X_n > x\sigma) = \Big(\Theta(x) + \theta\varepsilon_x\Big) \inf_{\lambda\geq 0} \mathbf{E}e^{\lambda(X_n - x\sigma)} \tag{12}$$

$$\leq \Big((\Theta(x) + \varepsilon_x) \wedge 1\Big) H_n(x,\sigma), \tag{13}$$

where

$$\varepsilon_x = \frac{e^t}{1-2t}\left(\frac{1.58}{\sqrt{\pi}}\frac{B}{\sigma} + \frac{2^{3+\delta}C_{2+\delta}}{(1-2t)^{\delta/2}} \frac{\sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}}{\sigma^{2+\delta}}\right)$$

with $t = \frac{2xB/\sigma}{1+\sqrt{1-4xB/\sigma}}$.

Notice that $\inf_{\lambda\geq 0}\mathbf{E}e^{\lambda(X_n - x\sigma)}$ is sometimes written in the form $\exp\{-n\Lambda_n^*(\frac{x\sigma}{n})\}$, where $\Lambda_n^*(x) = \sup_{\lambda\geq 0}\{\lambda x - \frac{1}{n}\log \mathbf{E}e^{\lambda X_n}\}$ is the Fenchel-Legendre transform of the cumulant function of X_n. When $(\xi_i)_{i=1,\dots,n}$ are i.i.d, we call $\Lambda^*(x) = \Lambda_n^*(x)$ the good rate function of LDP theory (see Deuschel and Stroock [11] or Dembo and Zeitouni [10]).

For a random variable ξ_i without moment larger than 2, some improvements of Hoeffding's inequality (2) can be find, for instance, in Eaton [13], Bentkus [3,4] and Pinelis [35]. In these paper, some larger classes of functions are considered instead of the class of exponential functions usually used in Markov's inequality. In particular, when ξ_i are bounded $\xi_i \leq 1$, Bentkus [4] showed that

$$\mathbf{P}\left(\sum_{i=1}^n \xi_i \geq x\right) \leq \frac{e^2}{2}\,\mathbf{P}^o\left(\sum_{i=1}^n \eta_i \geq x\right), \tag{14}$$

where η_i are i.i.d. with distribution (5), $\mathbf{P}^o\left(\sum_{i=1}^n \eta_i \geq x\right)$ is the log-concave hull of $\mathbf{P}(\sum_{i=1}^n \eta_i \geq x)$ and $\mathbf{P}^o \geq \mathbf{P}$. As $\eta_i \leq 1$, inequality (14) is sharp up to an absolute constant $\frac{e^2}{2}$. Here we give an equivalent result to (14). Applying (13) to (14) with $B = \max\{1, \frac{\sigma^2}{n}\}$, we find that if $\xi_i \leq 1$, then, for all $0 \leq x \leq 0.24\min\{\frac{n}{\sigma}, \sigma\}$,

$$\begin{aligned}\mathbf{P}(X_n \geq x\sigma) &\leq \frac{e^2}{2}\mathbf{P}^o\left(\sum_{i=1}^n \eta_i \geq x\sigma\right) \\ &= \frac{e^2}{2}\left(\Theta(x) + O(1)\max\left\{\frac{1}{\sigma}, \frac{\sigma}{n}\right\}\right) H_n(x,\,\sigma). \end{aligned} \tag{15}$$

This equivalent result (15) shows us that (14) improves Hoeffding's bound $H_n(x,\, \sigma)$ by adding a factor $F_n(x, \sigma) =: \frac{e^2}{2}\left(\varTheta(x) + O(1) \max\left\{\frac{1}{\sigma}, \frac{\sigma}{n}\right\}\right)$ in the range $0 \leq x \leq 0.24 \min\{\frac{n}{\sigma}, \sigma\}$. Moreover, in the i.i.d. case, we find that bound (14) converges to $\frac{e^2}{2}\varTheta(x) H_n(x,\, \sigma)$ for $0 \leq x = o(n^{\delta/2})$ as $n \to \infty$, which is worse than $\varTheta(x) H_n(x,\, \sigma)$ related to bound (13).

Using Theorem 1 and $C_3 \leq 0.56$, we easily obtain the following corollary.

Corollary 1 *Assume that $(\xi_i)_{i=1,...,n}$ satisfy $\xi_i \leq 1$ and $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ for some constant $B > 0$ and all i. Then, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$,*

$$\mathbf{P}(X_n > x\sigma) = \left(\varTheta(x) + 16\theta\frac{B}{\sigma}\right) \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)} \tag{16}$$

$$\leq \left(\left(\varTheta(x) + 16\frac{B}{\sigma}\right) \wedge 1\right) H_n(x, \sigma). \tag{17}$$

Note that the lees-than sign in Talagrand's inequality (6) is replaced by the equal sign in (16). Thus equality (16) completes Talagrand's inequality (6) by giving a lower bound. Since $\inf_{\lambda \geq 0} \mathbf{E} \exp\{\lambda(\sum_i^n \eta_i - x\sigma)\} = H_n(x, \sigma)$, by (16), we find that (17) attains to equality under the law (5). Hence Talagrand's inequality (17) is sharp regardless the constant 16.

Inequality (17) implies the following Cramér-type large deviations.

Corollary 2 *Assume that $(\xi_i)_{i=1,...,n}$ satisfy $\xi_i \leq 1$ and $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ for some constant $B > 0$ and all i. Then, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$,*

$$\mathbf{P}(X_n > x\sigma) \leq \Big(1 - \varPhi\left(\hat{x}\right)\Big) \left[1 + 16\sqrt{2\pi}\left(1 + \hat{x}\right)\frac{B}{\sigma}\right], \tag{18}$$

where $\hat{x} = \frac{x}{\sqrt{1 + \frac{x}{3\sigma}}}$ and satisfies

$$\hat{x} = x\left(1 - \frac{x}{6\sigma} + o(\frac{x}{\sigma})\right) \qquad \text{as} \qquad \frac{x}{\sigma} \to 0.$$

The interesting feature of the bound (18) is that it closely recovers the shape of the standard normal tail for all $0 \leq x = o(\sigma)$ as $\sigma \to \infty$.

The well-known asymptotic expansions of tail probabilities (see Petrov [31]) show that $\hat{x}$ in Corollary 2 usually can not be replaced by x. However, in the following sub-Gaussian case, $\hat{x}$ can be replaced by x.

Theorem 2 *Assume that $(\xi_i)_{i=1,...,n}$ satisfy $\xi_i \leq \sigma_i$ and $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ for some constant $B > 0$ and all i. Then, for all $0 \leq x < 0.25\frac{\sigma}{B}$,*

$$\mathbf{P}(X_n > x\sigma) \leq \Big(1 - \varPhi(x)\Big) \left[1 + c_x\left(1 + x\right)\frac{B}{\sigma}\right], \tag{19}$$

where

$$c_x = \frac{e^{t + t^2}}{1 - t}\left(\sqrt{2} + \frac{16\sqrt{2\pi}C_3 e^{\frac{1}{2}t^2}}{(1 - t)^{\frac{1}{2}}}\right)$$

with $t = \frac{2xB/\sigma}{1 + \sqrt{1 - 4xB/\sigma}}$. In particular, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$, we have $c_x \leq 32.47$.

Note that the condition $\xi_i \leq \sigma_i$ which replace the condition $\xi_i \leq 1$ in Corollary 2 is satisfied for Rademacher random variables (i.e., $\mathbf{P}(\xi_i = \pm 1) = \frac{1}{2}$).

For two-sided bounded random variables $|\xi_i| \leq B$ with $B > 0$, for example Rademacher random variables, the following theorem shows that the constants c_x in Theorem 1 can be further improved to a smaller one. Without loss of generality, we take $B = 1$, otherwise we consider ξ_i/B.

Theorem 3 *Assume that* $|\xi_i| \leq 1$ *for all* i*. Then, for all* $0 \leq x \leq 0.606\sigma$*,*

$$\mathbf{P}(X_n > x\sigma) = \Big(\varTheta(x) + \theta \frac{c_x}{\sigma}\Big) \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}, \tag{20}$$

where

$$c_x = \frac{2.24 e^{\frac{t^2}{2}}}{\sqrt{1-t}} + \frac{e^{t+t^2}}{\sqrt{\pi}(1-t)}$$

with $t = \frac{x}{\sigma} \exp\left\{\frac{ex^2}{2\sigma^2}\right\}$*. In particular, for all* $x \geq 0$*,*

$$\mathbf{P}(X_n > x\sigma) \leq \bigg(\Big(\varTheta(x) + \frac{c_x}{\sigma}\Big) \wedge 1\bigg) H_n(x, \sigma). \tag{21}$$

Moreover, if $0 \leq x \leq 0.1\sigma$*, then* $c_x \leq 3.08$*.*

It is clear that (21) improves the Hoeffding bound $H_n(x, \sigma)$ by adding a missing factor $(\varTheta(x) + c_x/\sigma) \wedge 1$.

3 Auxiliary Results

Assume that $\xi_i \leq B$ for some constant $B > 0$ and all i. We consider the positive random variable

$$Z_n(\lambda) = \prod_{i=1}^{n} \frac{e^{\lambda \xi_i}}{\mathbf{E} e^{\lambda \xi_i}}, \qquad \lambda \geq 0,$$

so that $\mathbf{E} Z_n(\lambda) = 1$ (the Esscher transformation). Introduce the *conjugate probability measure* $\mathbf{P}_\lambda$ defined by

$$d\mathbf{P}_\lambda = Z_n(\lambda) d\mathbf{P}. \tag{22}$$

Denote by $\mathbf{E}_\lambda$ the expectation with respect to $\mathbf{P}_\lambda$. Setting

$$b_i(\lambda) = \mathbf{E}_\lambda \xi_i = \frac{\mathbf{E} \xi_i e^{\lambda \xi_i}}{\mathbf{E} e^{\lambda \xi_i}}, \quad i = 1, ..., n,$$

and

$$\eta_i(\lambda) = \xi_i - b_i(\lambda), \quad i = 1, ..., n,$$

we obtain the semimartingale decomposition:

$$X_k = B_k(\lambda) + Y_k(\lambda), \quad k = 1, ..., n, \tag{23}$$

where

$$B_k(\lambda) = \sum_{i=1}^{k} b_i(\lambda) \qquad \text{and} \qquad Y_k(\lambda) = \sum_{i=1}^{k} \eta_i(\lambda).$$

In the proofs of Theorems 1-3, we shall need a two-sided bound of $B_n(\lambda)$. To this end, we need some technical lemmas.

For a random variable bounded from above, the following inequality is well-known.

Lemma 1 *Assume that* $\mathbf{E}\xi_i = 0$ *and* $\xi_i \leq 1$. *Then, for all* $\lambda \geq 0$,

$$\mathbf{E}e^{\lambda \xi_i} \leq Be(\lambda, \sigma_i^2) =: \frac{\sigma_i^2}{1+\sigma_i^2} \exp\{\lambda\} + \frac{1}{1+\sigma_i^2} \exp\left\{-\lambda \sigma_i^2\right\}.$$

A proof of this inequality can be found in Bennett [2]. The last inequality is sharp and attains to equality when

$$\mathbf{P}(\xi_i = 1) = \frac{\sigma_i^2}{1+\sigma_i^2} \quad \text{and} \quad \mathbf{P}(\xi_i = -\sigma_i^2) = \frac{1}{1+\sigma_i^2}.$$

By Lemma 1, we easily obtain the following estimation of the moment generating function $\mathbf{E}e^{\lambda \xi_i}$.

Lemma 2 *Assume that* $(\xi_i)_{i=1,...,n}$ *satisfy* $\xi_i \leq B$ *and* $\mathbf{E}|\xi_i|^{2+\delta} \leq B^{2+\delta}$ *for some constants* $B, \delta > 0$ *and all* i. *Then,* $\sigma_i^2 \leq B^2$ *and, for all* $\lambda \geq 0$,

$$\mathbf{E}e^{\lambda \xi_i} \leq e^{\frac{B^2\lambda^2}{2}}. \tag{24}$$

Proof. Using Jensen's inequality and $\mathbf{E}|\xi_i|^{2+\delta} \leq B^{2+\delta}$, we deduce

$$\sigma_i^2 = \mathbf{E}\xi_i^2 \leq (\mathbf{E}|\xi_i|^{2+\delta})^{2/(2+\delta)} \leq B^2.$$

Since $\frac{d}{dt}Be(\lambda, t) \geq 0$, for all $\lambda, t \geq 0$, the function $Be(\lambda, t)$ is increasing in $t \geq 0$ for all $\lambda \geq 0$. Hence, by Lemma 1 and $\frac{\sigma_i}{B} \leq 1$, for all $\lambda \geq 0$,

$$\mathbf{E}e^{\lambda \xi_i} = \mathbf{E}e^{\lambda B \frac{\xi_i}{B}} \leq \frac{e^{B\lambda} + e^{-B\lambda}}{2} \leq e^{\frac{B^2\lambda^2}{2}}. \tag{25}$$

This completes the proof of Lemma 2. □

In the following lemma, we give a two-sided bound for $B_n(\lambda)$.

Lemma 3 *Assume that* $\xi_i \leq B$ *for some constant* $B > 0$ *and all* i. *Then, for all* $\lambda \geq 0$,

$$B_n(\lambda) \leq \frac{e^{B\lambda} - 1}{B} \sigma^2.$$

If $(\xi_i)_{i=1,...,n}$ *satisfy condition (A) for all* i, *then, for all* $\lambda \geq 0$,

$$B_n(\lambda) \geq \left(1 - \frac{B\lambda}{2}\right) \lambda e^{-\frac{B^2\lambda^2}{2}} \sigma^2.$$

Proof. By Jensen's inequality, we have, for all $\lambda \geq 0$, $\mathbf{E}e^{\lambda\xi_i} \geq e^{\lambda \mathbf{E}\xi_i} = 1$. Since $\mathbf{E}\xi_i e^{\lambda\xi_i} = \mathbf{E}\xi_i(e^{\lambda\xi_i} - 1) \geq 0$, $\lambda \geq 0$, we obtain the first assertion, for all $\lambda \geq 0$,

$$\begin{aligned} B_n(\lambda) &\leq \sum_{i=1}^n \mathbf{E}\xi_i e^{\lambda\xi_i} = \sum_{i=1}^n \int_0^\lambda \mathbf{E}\xi_i^2 e^{t\xi_i}\, dt \\ &\leq \sum_{i=1}^n \int_0^\lambda \sigma_i^2\, e^{Bt} dt \\ &= \frac{e^{B\lambda} - 1}{B}\, \sigma^2. \end{aligned}$$

If $(\xi_i)_{i=1,...,n}$ satisfy condition (A), it follows that, for all $\lambda \geq 0$,

$$\begin{aligned}\sum_{i=1}^n \mathbf{E}\xi_i e^{\lambda \xi_i} &= \int_0^\lambda \sum_{i=1}^n \mathbf{E}\xi_i^2 e^{t\xi_i} dt \qquad (26)\\ &\geq \int_0^\lambda (1-Bt)dt \sum_{i=1}^n \mathbf{E}\xi_i^2 \\ &= \left(1-\frac{B\lambda}{2}\right)\lambda\sigma^2.\end{aligned}$$

Using Lemma 2, we get the lower bound of $B_n(\lambda)$, for all $\lambda \geq 0$,

$$B_n(\lambda) = \sum_{i=1}^n \frac{\mathbf{E}\xi_i e^{\lambda\xi_i}}{\mathbf{E}e^{\lambda\xi_i}} \geq \left(1-\frac{B\lambda}{2}\right)\lambda e^{-\frac{B^2\lambda^2}{2}}\sigma^2,$$

which completes the proof of Lemma 3. □

Write the *cumulant* function

$$\Psi_n(\lambda) = \sum_{i=1}^n \log \mathbf{E}e^{\lambda\xi_i}, \qquad \lambda \geq 0. \qquad (27)$$

We can easily obtain the following upper bound for the function $\Psi_n(\lambda)$.

Lemma 4 *Assume that* $\xi_i \leq 1$ *for all* i. *Then, for all* $\lambda \geq 0$,

$$\Psi_n(\lambda) \leq n \log\left(\frac{1}{1+\sigma^2/n}\exp\left\{-\lambda\sigma^2/n\right\} + \frac{\sigma^2/n}{1+\sigma^2/n}\exp\{\lambda\}\right).$$

Proof. Since the function

$$f(\lambda, t) = \log\left(\frac{1}{1+t}\exp\left\{-\lambda t\right\} + \frac{t}{1+t}\exp\{\lambda\}\right), \qquad \lambda, t \geq 0,$$

has a negative second derivative for $t > 0$ (see Lemma 3 in Hoeffding [20]), then, for any fixed $\lambda \geq 0$, $-f(\lambda, t)$ is convex in $t \geq 0$ and

$$f(\lambda, t) \leq f(\lambda, 0) + \frac{\partial}{\partial t} f(\lambda, 0)\, t = (e^\lambda - 1 - \lambda)\, t, \qquad t \geq 0. \qquad (28)$$

Therefore by Lemma 1 and Jensen's inequality, we get, for all $\lambda \geq 0$,

$$\begin{aligned}\Psi_n(\lambda) &\leq \sum_{i=1}^n f(\lambda, \sigma_i^2) \\ &\leq n f(\lambda, \sigma^2/n) \\ &= n\log\left(\frac{1}{1+\sigma^2/n}\exp\left\{-\lambda\sigma^2/n\right\} + \frac{\sigma^2/n}{1+\sigma^2/n}\exp\{\lambda\}\right).\end{aligned}$$

This completes the proof of Lemma 4. □

For random variable $Y_n(\lambda)$, $\lambda \geq 0$, we have a rate of convergence to the standard normal law. Before present the result, we should firstly introduce into the conjugate variance of σ^2. Denote the variance of $Y_n(\lambda)$ by $\overline{\sigma}^2(\lambda) = \mathbf{E}_\lambda Y_n^2(\lambda),\ \lambda \geq 0$. By the relation between $\mathbf{E}$ and $\mathbf{E}_\lambda$, the following inequality is obvious

$$\overline{\sigma}^2(\lambda) = \sum_{i=1}^n \left(\frac{\mathbf{E}\xi_i^2 e^{\lambda\xi_i}}{\mathbf{E}e^{\lambda\xi_i}} - \frac{(\mathbf{E}\xi_i e^{\lambda\xi_i})^2}{(\mathbf{E}e^{\lambda\xi_i})^2}\right), \quad \lambda \geq 0.$$

The following lemma gives some estimations of $\overline{\sigma}^2(\lambda)$.

Lemma 5 *Assume that* $(\xi_i)_{i=1,\ldots,n}$ *satisfy* $\xi_i \leq B$ *and condition (A) for all* i*. Then, for all* $\lambda \geq 0$,

$$(1-2B\lambda)^+\sigma^2 \leq \overline{\sigma}^2(\lambda) \leq e^{B\lambda}\sigma^2. \tag{29}$$

Proof. Since $\mathbf{E}e^{\lambda\xi_i} \geq 1, \lambda \geq 0$, and $\xi_i \leq B$, we get, for all $\lambda \geq 0$,

$$\overline{\sigma}^2(\lambda) \leq \sum_{i=1}^n \mathbf{E}\xi_i^2 e^{\lambda\xi_i} \leq \sum_{i=1}^n \mathbf{E}\xi_i^2 e^{\lambda B} = e^{B\lambda}\sigma^2.$$

This gives the upper bound of $\overline{\sigma}^2(\lambda)$. For all $\lambda \geq 0$, it is easy to see that

$$\mathbf{E}\xi_i e^{\lambda\xi_i} = \int_0^\lambda \mathbf{E}\xi_i^2 e^{t\xi_i}dt \leq \int_0^\lambda e^{Bt}dt\mathbf{E}\xi_i^2 = \left(\frac{e^{B\lambda}-1}{B}\right)\mathbf{E}\xi_i^2. \tag{30}$$

Using Lemma 2 and condition (A), we obtain, for all $\lambda \geq 0$,

$$\begin{aligned}\overline{\sigma}^2(\lambda) &\geq \sum_{i=1}^n \frac{\mathbf{E}\xi_i^2 e^{\lambda\xi_i} - (\mathbf{E}\xi_i e^{\lambda\xi_i})^2}{e^{B^2\lambda^2}} \\ &\geq \frac{(1-B\lambda)\sigma^2 - (e^{B\lambda}-1)^2 B^{-2}\sum_{i=1}^n(\mathbf{E}\xi_i^2)^2}{e^{B^2\lambda^2}} \\ &\geq e^{-B^2\lambda^2}\left(1-B\lambda-(e^{B\lambda}-1)^2\right)\sigma^2.\end{aligned}$$

Noting that $\overline{\sigma}^2(\lambda) \geq 0$, by a simple calculation, we have, for all $\lambda \geq 0$,

$$\overline{\sigma}^2(\lambda) \geq (1-2B\lambda)^+\sigma^2,$$

which gives the lower bound of $\overline{\sigma}^2(\lambda)$. □

Next, we present the rate of convergence of $Y_n(\lambda)$ to the standard normal law.

Lemma 6 *Assume that* $(\xi_i)_{i=1,\ldots,n}$ *satisfy* $\xi_i \leq B$ *and* $\mathbf{E}|\xi_i|^{2+\delta} < \infty$ *for some constants* $\delta \in (0,1]$, $B > 0$ *and all* i*. Then, for all* $\lambda \geq 0$,

$$\sup_{y\in\mathbf{R}}\left|\mathbf{P}_\lambda\left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y\right) - \Phi(y)\right| \leq \frac{2^{2+\delta}C_{2+\delta}e^{B\lambda}}{\overline{\sigma}^{2+\delta}(\lambda)}\sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}.$$

Proof. Notice that $Y_n(\lambda) = \sum_{i=1}^n \eta_i(\lambda)$ is the sum of independent random variables $\eta_i(\lambda)$ and $\mathbf{E}_\lambda\eta_i(\lambda) = 0$. Using the well-known rate of convergence in the central limit theorem (cf. e.g. Petrov [31], p. 115), we get, for all $\lambda \geq 0$,

$$\sup_{y\in\mathbf{R}}\left|\mathbf{P}_\lambda\left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y\right) - \Phi(y)\right| \leq \frac{C_{2+\delta}}{\overline{\sigma}^{2+\delta}(\lambda)}\sum_{i=1}^n \mathbf{E}_\lambda|\eta_i|^{2+\delta}.$$

Using the inequality $(a+b)^{1+q} \leq 2^q(a^{1+q}+b^{1+q})$ for $a, b, q \geq 0$, we deduce, for all $\lambda \geq 0$,

$$\begin{aligned}\sum_{i=1}^n \mathbf{E}_\lambda|\eta_i|^{2+\delta} &\leq 2^{1+\delta}\sum_{i=1}^n \mathbf{E}_\lambda(|\xi_i|^{2+\delta} + |\mathbf{E}_\lambda\xi_i|^{2+\delta}) \\ &\leq 2^{2+\delta}\sum_{i=1}^n \mathbf{E}_\lambda|\xi_i|^{2+\delta} \\ &\leq 2^{2+\delta}\sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}e^{\lambda\xi_i} \\ &\leq 2^{2+\delta}e^{B\lambda}\sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}.\end{aligned}$$

Therefore, we obtain, for all $\lambda \geq 0$,

$$\sup_{y \in \mathbf{R}} \left| \mathbf{P}_\lambda \left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y \right) - \Phi(y) \right| \leq \frac{2^{2+\delta} C_{2+\delta} e^{B\lambda}}{\overline{\sigma}^{2+\delta}(\lambda)} \sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}.$$

This completes the proof of Lemma 6. □

We are now ready to prove our second main result.

Theorem 4 *Assume that* $(\xi_i)_{i=1,\ldots,n}$ *satisfy* $\xi_i \leq B$ *and* $\mathbf{E}|\xi_i|^{2+\delta} < \infty$ *for some constants* $\delta \in (0,1]$, $B > 0$ *and all* i. *Then, for all* $x \geq 0$,

$$\mathbf{P}(X_n > x\sigma) = \Big(\Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) + \theta \varepsilon_x \Big) \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}, \tag{31}$$

where

$$\varepsilon_x = \frac{2^{3+\delta} C_{2+\delta} e^{B\overline{\lambda}}}{\overline{\sigma}^{2+\delta}(\overline{\lambda})} \sum_{i=1}^n \mathbf{E}|\xi_i|^{2+\delta}$$

with $\overline{\lambda}$ *satisfying* $\Psi_n'(\overline{\lambda}) = x\sigma$.

The previous theorem is closely related to Talagrand's inequality (6) (cf. Theorem 3.3 in [37] for details). Here, we would like to highlight some differences between the two results. First, the less-than sign in Talagrand's inequality (6) has been completed to the equal sign in (31). Second, Talagrand's inequality (6) holds for $0 \leq x = O(\frac{\sigma}{B})$, while Theorem 4 holds for all $x \geq 0$.

Proof of Theorem 4. According to (22), we have the following representation: for all $x, \lambda \geq 0$,

$$\begin{aligned}
&\mathbf{P}(X_n > x\sigma) \\
&= \mathbf{E}_\lambda (Z_n(\lambda)^{-1} \mathbf{1}_{\{X_n > x\sigma\}}) \\
&= \mathbf{E}_\lambda (e^{-\lambda X_n + \Psi_n(\lambda)} \mathbf{1}_{\{X_n > x\sigma\}}) \\
&= \mathbf{E}_\lambda (e^{-\lambda x\sigma + \Psi_n(\lambda) - \lambda Y_n(\lambda) - \lambda B_n(\lambda) + \lambda x\sigma} \mathbf{1}_{\{Y_n(\lambda) + B_n(\lambda) - x\sigma > 0\}}) \\
&= e^{-\lambda x\sigma + \Psi_n(\lambda)} \mathbf{E}_\lambda (e^{-\lambda[Y_n(\lambda) + B_n(\lambda) - x\sigma]} \mathbf{1}_{\{Y_n(\lambda) + B_n(\lambda) - x\sigma > 0\}}).
\end{aligned}$$

Setting $U_n(\lambda) = \lambda(Y_n(\lambda) + B_n(\lambda) - x\sigma)$, we get

$$\mathbf{P}(X_n > x\sigma) = e^{-\lambda x\sigma + \Psi_n(\lambda)} \int_0^\infty e^{-t} \mathbf{P}_\lambda (0 < U_n(\lambda) \leq t) dt. \tag{32}$$

The exponential function $e^{-\lambda x\sigma + \Psi_n(\lambda)}$ in (32) attains its minimum at $\lambda = \overline{\lambda}$, where $\overline{\lambda}$ satisfies $\Psi_n'(\overline{\lambda}) = x\sigma$. Then, $B_n(\overline{\lambda}) = \Psi_n'(\overline{\lambda}) = x\sigma$, $U_n(\overline{\lambda}) = \overline{\lambda} Y_n(\overline{\lambda})$ and

$$e^{-\overline{\lambda} x\sigma + \Psi_n(\overline{\lambda})} = \inf_{\lambda \geq 0} e^{-\lambda x\sigma + \Psi_n(\lambda)} = \inf_{\lambda \geq 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}. \tag{33}$$

Using Lemma 6, we deduce

$$\begin{aligned}
&\int_0^\infty e^{-t} \mathbf{P}_{\overline{\lambda}} (0 < U_n(\overline{\lambda}) \leq t) dt \qquad (34) \\
&= \int_0^\infty e^{-\overline{\lambda} y \overline{\sigma}(\overline{\lambda})} \mathbf{P}_{\overline{\lambda}} \left(0 < U_n(\overline{\lambda}) \leq \overline{\lambda} y \overline{\sigma}(\overline{\lambda})\right) \overline{\lambda}\overline{\sigma}(\overline{\lambda}) dy \\
&= \int_0^\infty e^{-\overline{\lambda} y \overline{\sigma}(\overline{\lambda})} \mathbf{P}(0 < \mathcal{N}(0,1) \leq y) \overline{\lambda}\overline{\sigma}(\overline{\lambda}) dy + \theta \varepsilon_x \\
&= \int_0^\infty e^{-\overline{\lambda} y \overline{\sigma}(\overline{\lambda})} d\Phi(y) + \theta \varepsilon_x \\
&= \Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) + \theta \varepsilon_x,
\end{aligned}$$

where

$$\varepsilon_x = \frac{2^{3+\delta} C_{2+\delta} e^{B\overline{\lambda}}}{\overline{\sigma}^{2+\delta}(\overline{\lambda})} \sum_{i=1}^{n} \mathbf{E}|\xi_i|^{2+\delta}.$$

Therefore, from (32), for all $x \geq 0$,

$$\mathbf{P}(X_n > x\sigma) = \Big(\Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) + \theta\varepsilon_x\Big) \inf_{\lambda \geq 0} \mathbf{E}e^{\lambda(X_n - x\sigma)}.$$

This completes the proof of Theorem 4. $\square$

4 Proof of Theorem 1

In the spirit of Talagrand [37], we make use of $\Theta(x)$ to approximate $\Theta(\overline{\lambda}\overline{\sigma}(\overline{\lambda}))$ of Theorem 4. The following proof of Theorem 1 is a continuation of the proof of Theorem 4.

Proof of Theorem 1. Using (33) and Lemma 4, we get, for all $x \geq 0$,

$$\begin{aligned} &\inf_{\lambda \geq 0} \mathbf{E}e^{\lambda(X_n - x\sigma)} \\ &\leq \inf_{\lambda \geq 0} \exp\left\{-\lambda x\sigma + n\log\left(\frac{1}{1+\sigma^2/n}\exp\left\{-\lambda\sigma^2/n\right\} + \frac{\sigma^2/n}{1+\sigma^2/n}\exp\{\lambda\}\right)\right\} \\ &= H_n(x, \sigma). \end{aligned} \tag{35}$$

Since $|\Theta'(x)| \leq \frac{1}{\sqrt{\pi}x^2}$, we deduce

$$\left|\Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) - \Theta(x)\right| \leq \frac{1}{\sqrt{\pi}} \frac{|x - \overline{\lambda}\overline{\sigma}(\overline{\lambda})|}{\overline{\lambda}^2\overline{\sigma}^2(\overline{\lambda}) \wedge x^2}. \tag{36}$$

Using Lemma 3 and the inequality $e^{-x} \geq 1 - x$, $x \geq 0$, we have, for all $0 \leq \overline{\lambda} \leq \frac{1}{B}$,

$$\left(1 - B\overline{\lambda}\right)\overline{\lambda}\sigma \leq \left(1 - \frac{B\overline{\lambda}}{2}\right)\overline{\lambda}e^{-\frac{B^2\overline{\lambda}^2}{2}}\sigma \leq \frac{B_n(\overline{\lambda})}{\sigma} = x \leq \frac{e^{B\overline{\lambda}} - 1}{B}\sigma. \tag{37}$$

By the estimation of $\overline{\sigma}(\overline{\lambda})$ in Lemma 5, it follows that, for all $0 \leq \overline{\lambda} \leq \frac{1}{2B}$,

$$\begin{aligned} &\left|x - \overline{\lambda}\overline{\sigma}(\overline{\lambda})\right| \\ &\leq \overline{\lambda}\sigma\left[\left(\frac{e^{B\overline{\lambda}} - 1}{B\overline{\lambda}} - \sqrt{1 - 2B\overline{\lambda}}\right) \vee \left(e^{\frac{B\overline{\lambda}}{2}} - (1 - B\overline{\lambda})\right)\right] \\ &\leq 1.58 e^{B\overline{\lambda}} B\overline{\lambda}^2\sigma \end{aligned} \tag{38}$$

and

$$\begin{aligned} \overline{\lambda}^2\overline{\sigma}^2(\overline{\lambda}) \wedge x^2 &\geq \left(\overline{\lambda}^2\sigma^2(1 - 2B\overline{\lambda})\right) \wedge \left(\overline{\lambda}^2\sigma^2(1 - B\overline{\lambda})^2\right) \\ &= \overline{\lambda}^2\sigma^2(1 - 2B\overline{\lambda}). \end{aligned} \tag{39}$$

Hence, (36) implies that, for all $0 \leq \overline{\lambda} < \frac{1}{2B}$,

$$\left|\Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) - \Theta(x)\right| \leq \frac{1.58}{\sqrt{\pi}} \frac{e^{B\overline{\lambda}}}{(1 - 2B\overline{\lambda})} \frac{B}{\sigma}. \tag{40}$$

By (37), it follows that $\left(1-B\overline{\lambda}\right)\overline{\lambda} \leq \frac{x}{\sigma}$ and

$$0 \leq B\overline{\lambda} \leq t =: \frac{2xB/\sigma}{1+\sqrt{1-4Bx/\sigma}} \quad \text{for all} \quad 0 \leq x\frac{B}{\sigma} < 0.25. \tag{41}$$

Therefore, by Lemma 5, it is easy to see that

$$\frac{e^{B\overline{\lambda}}}{\overline{\sigma}^{2+\delta}(\overline{\lambda})}\sum_{i=1}^{n}\mathbf{E}|\xi_i|^{2+\delta} \leq \frac{e^{t}}{(1-2t)^{1+\frac{\delta}{2}}}\frac{\sum_{i=1}^{n}\mathbf{E}|\xi_i|^{2+\delta}}{\sigma^{2+\delta}}. \tag{42}$$

Combining (31) and (40)-(42) together, we have, for all $0 \leq x < 0.25\frac{\sigma}{B}$,

$$\begin{aligned}\mathbf{P}(X_n > x\sigma) &= \left(\Theta(x) + \theta\varepsilon_x\right)\inf_{\lambda\geq 0}\mathbf{E}e^{\lambda(X_n - x\sigma)} \\ &\leq \left((\Theta(x)+\varepsilon_x)\wedge 1\right)H_n(x,\sigma),\end{aligned}$$

where

$$\varepsilon_x = \frac{e^t}{1-2t}\left(\frac{1.58}{\sqrt{\pi}}\frac{B}{\sigma} + \frac{2^{3+\delta}C_{2+\delta}}{(1-2t)^{\delta/2}}\frac{\sum_{i=1}^{n}\mathbf{E}|\xi_i|^{2+\delta}}{\sigma^{2+\delta}}\right).$$

This completes the proof of Theorem 1. □

Proof of Corollary 2. Since $\Theta(x)$ is decreasing in $x \geq 0$, we deduce $\Theta(x) \leq \Theta(\hat{x})$ where $\hat{x} = \frac{x}{\sqrt{1+\frac{x}{3\sigma}}}$. Notice that Hoeffding's bound is less than Bernstein's bound, i.e.

$$H_n(x,\sigma) \leq \exp\left\{-\frac{\widehat{x}^2}{2}\right\}$$

(cf. Remark 2.1 of [14]). Therefore, from (17), we have, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$,

$$\begin{aligned}\mathbf{P}(X_n > x\sigma) &\leq \left(\Theta(\hat{x}) + 16\frac{B}{\sigma}\right)\exp\left\{-\frac{\hat{x}^2}{2}\right\} \\ &= 1 - \Phi(\hat{x}) + 16\frac{B}{\sigma}\exp\left\{-\frac{\hat{x}^2}{2}\right\}.\end{aligned}$$

Using (8), we obtain, for all $0 \leq x \leq 0.1\frac{\sigma}{B}$,

$$\mathbf{P}(X_n > x\sigma) \leq (1-\Phi(\hat{x}))\left[1 + 16\sqrt{2\pi}\,(1+\hat{x})\,\frac{B}{\sigma}\right].$$

This completes the proof of Corollary 2. □

5 Proof of Theorem 2

Under the condition of Theorem 2, we have the following upper bound of $\Psi_n(\lambda)$.

Lemma 7 *Assume that* $\xi_i \leq \sigma_i$ *for all* i*. Then, for all* $\lambda \geq 0$,

$$\Psi_n(\lambda) \leq \frac{\lambda^2\sigma^2}{2}.$$

Proof. Using Lemma 1, we have, for all $t \geq 0$,

$$\mathbf{E}\exp\left\{t\frac{\xi_i}{\sigma_i}\right\} \leq \frac{e^{-t}+e^{t}}{2} \leq \exp\left\{\frac{t^2}{2}\right\}.$$

Hence, for all $\lambda \geq 0$,

$$\mathbf{E}e^{\lambda\xi_i} \leq \exp\left\{\frac{\lambda^2\sigma_i^2}{2}\right\} \qquad \text{and} \qquad \Psi_n(\lambda) \leq \frac{\lambda^2}{2}\sum_{i=1}^{n}\sigma_i^2 = \frac{\lambda^2\sigma^2}{2}.$$

This gives the upper bound of $\Psi_n(\lambda)$. □

Under the condition of Theorem 2, we have the following lower bound of $\overline{\sigma}^2(\lambda)$.

Lemma 8 *Assume that* $\xi_i \leq B$ *and* $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ *for some constant* $B > 0$ *and all* i*. Then, for all* $\lambda \geq 0$,

$$(1-B\lambda)^+ e^{-B^2\lambda^2}\sigma^2 \leq \overline{\sigma}^2(\lambda). \tag{43}$$

Proof. Denote by $f(\lambda) = \mathbf{E}\xi_i^2 e^{\lambda\xi_i}\mathbf{E}e^{\lambda\xi_i} - (\mathbf{E}\xi_i e^{\lambda\xi_i})^2$. It is easy to see that

$$f'(0) = \mathbf{E}\xi_i^3 \quad \text{and} \quad f''(\lambda) = \mathbf{E}\xi_i^4 e^{\lambda\xi_i}\mathbf{E}e^{\lambda\xi_i} - (\mathbf{E}\xi_i^2 e^{\lambda\xi_i})^2 \geq 0.$$

Thus, for all $\lambda \geq 0$,

$$f(\lambda) \geq f(0) + f'(0)\lambda = \mathbf{E}\xi_i^2 + \lambda\mathbf{E}\xi_i^3. \tag{44}$$

Therefore, for all $\lambda \geq 0$,

$$\mathbf{E}_\lambda\eta_i^2 = \frac{\mathbf{E}\xi_i^2 e^{\lambda\xi_i}\mathbf{E}e^{\lambda\xi_i} - (\mathbf{E}\xi_i e^{\lambda\xi_i})^2}{(\mathbf{E}e^{\lambda\xi_i})^2} \geq \frac{\mathbf{E}\xi_i^2 + \lambda\mathbf{E}\xi_i^3}{(\mathbf{E}e^{\lambda\xi_i})^2}.$$

Using $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$, we get, for all $\lambda \geq 0$,

$$\mathbf{E}_\lambda\eta_i^2 \geq \frac{\mathbf{E}\xi_i^2 + \lambda\mathbf{E}\xi_i^3}{(\mathbf{E}e^{\lambda\xi_i})^2} \geq \frac{(1-B\lambda)\mathbf{E}\xi_i^2}{(\mathbf{E}e^{\lambda\xi_i})^2}. \tag{45}$$

Since $\mathbf{E}|\xi_i|^3 \leq B\mathbf{E}\xi_i^2$ implies that $\mathbf{E}|\xi_i|^3 \leq B^3$, by Lemma 2 and $\overline{\sigma}^2(\lambda) \geq 0$, it follows that

$$\overline{\sigma}^2(\lambda) \geq \sum_{i=1}^{n}\frac{(1-B\lambda)^+\mathbf{E}\xi_i^2}{e^{B^2\lambda^2}} \geq (1-B\lambda)^+ e^{-B^2\lambda^2}\sigma^2.$$

This completes the proof of Lemma 8. □

Notice that the condition in Theorem 2 is stronger than that in Theorem 1. We can easily prove Theorem 2 by (12) in Theorem 1. However, in order to obtain a constant c_x in Theorem 2 as small as possible, we make use of Theorem 4 to prove Theorem 2.

Proof of Theorem 2. By Lemma 7, for all $x \geq 0$, we have

$$\begin{aligned}\inf_{\lambda\geq 0} \mathbf{E}e^{\lambda(X_n - x\sigma)} &= \inf_{\lambda\geq 0} \exp\left\{-\lambda x\sigma + \varPsi_n(\lambda)\right\} \\ &\leq \inf_{\lambda\geq 0} \exp\left\{-\lambda x\sigma + \frac{\lambda^2\sigma^2}{2}\right\} \\ &\leq \exp\left\{-\frac{x^2}{2}\right\}. \qquad (46)\end{aligned}$$

Since $\sigma_i^3 \leq \mathbf{E}|\xi_i|^3 \leq B\sigma_i^2$ and $\xi_i \leq \sigma_i$, we have $\sigma_i \leq B$ and $\xi_i \leq B$. Then, by Lemmas 5, 8 and inequality (37), it follows that, for all $0 \leq \overline{\lambda} < \frac{1}{B}$,

$$\begin{aligned}&\left|x - \overline{\lambda}\overline{\sigma}(\overline{\lambda})\right| \\ &\leq \overline{\lambda}\sigma\left[\left(\frac{e^{B\overline{\lambda}}-1}{B\overline{\lambda}} - \sqrt{1-B\overline{\lambda}}e^{-\frac{B^2\overline{\lambda}^2}{2}}\right) \vee \left(e^{\frac{B\overline{\lambda}}{2}} - \left(1-\frac{B\overline{\lambda}}{2}\right)e^{-\frac{B^2\overline{\lambda}^2}{2}}\right)\right] \\ &\leq e^{B\overline{\lambda}}B\overline{\lambda}^2\sigma \qquad (47)\end{aligned}$$

and

$$\begin{aligned}\overline{\lambda}^2\overline{\sigma}^2(\overline{\lambda}) \wedge x^2 &\geq \left(\overline{\lambda}^2\sigma^2(1-B\overline{\lambda})e^{-\frac{B^2\overline{\lambda}^2}{2}}\right) \wedge \left(\overline{\lambda}^2\sigma^2\left(1-\frac{B\overline{\lambda}}{2}\right)^2 e^{-B^2\overline{\lambda}^2}\right) \\ &\geq \overline{\lambda}^2\sigma^2(1-B\overline{\lambda})^{1.3}. \qquad (48)\end{aligned}$$

Hence, by (36), we obtain, for all $0 \leq \overline{\lambda} < \frac{1}{B}$,

$$\left|\varTheta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) - \varTheta(x)\right| \leq \frac{e^{B\overline{\lambda}}}{\sqrt{\pi}(1-B\overline{\lambda})^{1.3}}\frac{B}{\sigma}. \qquad (49)$$

By Lemma 8, it is easy to see that, for all $0 \leq \overline{\lambda} < \frac{1}{B}$,

$$\frac{e^{B\overline{\lambda}}}{\overline{\sigma}^3(\overline{\lambda})}\sum_{i=1}^n \mathbf{E}|\xi_i|^3 \leq \frac{e^{B\overline{\lambda}+\frac{3}{2}B^2\overline{\lambda}^2}}{(1-B\overline{\lambda})^{\frac{3}{2}}}\frac{B}{\sigma}.$$

Thus, from Theorem 4, it follows that, for all $0 \leq x < 0.25\frac{\sigma}{B}$,

$$\begin{aligned}\mathbf{P}(X_n > x\sigma) &\leq \left(\varTheta(x) + c_x\frac{B}{\sigma}\right)\exp\left\{-\frac{x^2}{2}\right\} \\ &= 1 - \varPhi(x) + c_x\frac{B}{\sigma}\exp\left\{-\frac{x^2}{2}\right\},\end{aligned}$$

where

$$c_x = \frac{e^{t+t^2}}{1-t}\left(\sqrt{2} + \frac{16\sqrt{2\pi}C_3e^{\frac{1}{2}t^2}}{(1-t)^{\frac{1}{2}}}\right)$$

with t defined in (41). Since $\varTheta(x) \geq \frac{1}{\sqrt{2\pi}(1+x)}, x \geq 0$, we have, for all $0 \leq x < 0.25\frac{\sigma}{B}$,

$$\frac{\mathbf{P}(X_n > x\sigma)}{1-\varPhi(x)} \leq 1 + \sqrt{2\pi}c_x(1+x)\frac{B}{\sigma}.$$

In particular, if $0 \leq x \leq 0.1\frac{\sigma}{B}$, using $C_3 \leq 0.56$, we have $\sqrt{2\pi}c_x \leq 32.47$. □

6 Proof of Theorem 3

For bounded random variables, the convergence rate of $Y_n(\lambda)$ to the standard normal law in Lemma 6 can be improved to the following one.

Lemma 9 *Assume that* $(\xi_i)_{i=1,\ldots,n}$ *satisfy* $|\xi_i| \leq 1$ *for all* i. *Then, for all* $\lambda \geq 0$,

$$\sup_{y\in\mathbf{R}} \left| \mathbf{P}_\lambda\left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y\right) - \varPhi(y) \right| \leq \frac{1.12}{\overline{\sigma}(\lambda)}.$$

Proof. Using the well-known rate of convergence in the central limit theorem in Petrov [31], p. 115, and $C_3 \leq 0.56$, we get, for all $\lambda \geq 0$,

$$\sup_{y\in\mathbf{R}} \left| \mathbf{P}_\lambda\left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y\right) - \varPhi(y) \right| \leq \frac{0.56}{\overline{\sigma}^3(\lambda)} \sum_{i=1}^n \mathbf{E}_\lambda|\eta_i|^3.$$

Since $|\eta_i| \leq |\xi_i| + \mathbf{E}_\lambda|\xi_i| \leq 2$, it follows that $\sum_{i=1}^n \mathbf{E}_\lambda|\eta_i|^3 \leq 2\overline{\sigma}^2(\lambda)$. Therefore, for all $\lambda \geq 0$,

$$\sup_{y\in\mathbf{R}} \left| \mathbf{P}_\lambda\left(\frac{Y_n(\lambda)}{\overline{\sigma}(\lambda)} \leq y\right) - \varPhi(y) \right| \leq \frac{1.12}{\overline{\sigma}(\lambda)},$$

which completes the proof of Lemma 9. ☐

Lemma 10 *Assume that* $(\xi_i)_{i=1,\ldots,n}$ *satisfy* $|\xi_i| \leq 1$ *for all* i. *Then, for all* $\lambda \geq 0$,

$$B_n(\lambda) \geq (1-e^{-\lambda})e^{-\frac{\lambda^2}{2}}\sigma^2.$$

Proof. Since $\xi_i \geq -1$, for all $\lambda \geq 0$,

$$\begin{aligned} \sum_{i=1}^n \mathbf{E}\xi_i e^{\lambda\xi_i} &= \sum_{i=1}^n \int_0^\lambda \mathbf{E}\xi_i^2 e^{t\xi_i}\,dt \\ &\geq \int_0^\lambda e^{-t}dt \sum_{i=1}^n \mathbf{E}\xi_i^2 \\ &= (1-e^{-\lambda})\sigma^2. \end{aligned} \tag{50}$$

Using Lemma 2 with $B = 1$, we get the lower bound of $B_n(\lambda)$: for all $\lambda \geq 0$,

$$B_n(\lambda) = \sum_{i=1}^n \frac{\mathbf{E}\xi_i e^{\lambda\xi_i}}{\mathbf{E}e^{\lambda\xi_i}} \geq (1-e^{-\lambda})e^{-\frac{\lambda^2}{2}}\sigma^2.$$

This completes the proof of Lemma 10. ☐

Proof of Theorem 3. The proof is close to the proof of Theorem 1, but here $B = 1$. From (34), by Lemma 9, we deduce

$$\begin{aligned} &\int_0^\infty e^{-t}\mathbf{P}_{\overline{\lambda}}(0 < U_n(\overline{\lambda}) \leq t)dt \\ &= \int_0^\infty e^{-\overline{\lambda}y\overline{\sigma}(\overline{\lambda})}\mathbf{P}_{\overline{\lambda}}\left(0 < U_n(\overline{\lambda}) \leq \overline{\lambda}y\sigma(\overline{\lambda})\right)\overline{\lambda}\sigma(\overline{\lambda})dy \\ &= \int_0^\infty e^{-\overline{\lambda}y\overline{\sigma}(\overline{\lambda})}\mathbf{P}\left(0 < \mathcal{N}(0,1) \leq y\right)\overline{\lambda}\sigma(\overline{\lambda})dy + \theta\frac{2.24}{\overline{\sigma}(\overline{\lambda})} \\ &= \int_0^\infty e^{-\overline{\lambda}y\overline{\sigma}(\overline{\lambda})}d\varPhi\left(y\right) + \theta\frac{2.24}{\overline{\sigma}(\overline{\lambda})} \\ &= \varTheta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) + \theta\frac{2.24}{\overline{\sigma}(\overline{\lambda})}, \end{aligned} \tag{51}$$

where $\overline{\lambda}$ satisfies $\Psi_n'(\overline{\lambda}) = B_n(\overline{\lambda}) = x\sigma$. Therefore, we obtain the corresponding result of Theorem 4 for two-sided bounded random variables: for all $x \ge 0$,

$$\mathbf{P}(X_n > x\sigma) = \left(\Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) + \theta \frac{2.24}{\overline{\sigma}(\overline{\lambda})} \right) \inf_{\lambda \ge 0} \mathbf{E} e^{\lambda(X_n - x\sigma)}. \tag{52}$$

Since $\overline{\lambda} e^{-\frac{\overline{\lambda}^2}{2}} \sigma^2 \le (1 - e^{-\overline{\lambda}}) e^{-\frac{\overline{\lambda}^2}{2}} \sigma^2 \le B_n(\overline{\lambda}) = x\sigma$ (cf. Lemma 10), by Lemma 8, we have, for all $0 \le \overline{\lambda} < 1$,

$$\overline{\lambda} \le \frac{x}{\sigma} e^{\frac{\overline{\lambda}^2}{2}} \le \frac{x}{\sigma} \exp\left\{ \frac{x^2 e^{\overline{\lambda}^2}}{2\sigma^2} \right\} \le t =: \frac{x}{\sigma} \exp\left\{ \frac{e\, x^2}{2\sigma^2} \right\}. \tag{53}$$

and

$$\begin{aligned} \overline{\lambda}^2 \overline{\sigma}^2(\overline{\lambda}) \wedge x^2 &\ge \overline{\lambda}^2 (1 - \overline{\lambda}) e^{-\overline{\lambda}^2} \sigma^2 \wedge \overline{\lambda}^2 e^{-\overline{\lambda}^2} \sigma^2 \\ &= \overline{\lambda}^2 (1 - \overline{\lambda}) e^{-\overline{\lambda}^2} \sigma^2. \end{aligned} \tag{54}$$

From (36) and (47), it follows that, for all $0 \le \overline{\lambda} < 1$,

$$\left| \Theta\left(\overline{\lambda}\overline{\sigma}(\overline{\lambda})\right) - \Theta(x) \right| \le \frac{e^{\overline{\lambda} + \overline{\lambda}^2}}{\sqrt{\pi}(1 - \overline{\lambda})} \frac{1}{\sigma}.$$

Using Lemma 8 again, we obtain, for all $0 \le \overline{\lambda} < 1$,

$$\frac{2.24}{\overline{\sigma}(\overline{\lambda})} \le \frac{2.24 e^{\frac{\overline{\lambda}^2}{2}}}{\sqrt{1 - \overline{\lambda}}} \frac{1}{\sigma}. \tag{55}$$

Returning to (52), we get, for all $0 \le \overline{\lambda} < 1$,

$$\mathbf{P}(X_n > x\sigma) = \left(\Theta(x) + \theta \frac{c_x(\overline{\lambda})}{\sigma} \right) \inf_{\lambda \ge 0} \mathbf{E} e^{\lambda(X_n - x\sigma)},$$

where

$$c_x(\overline{\lambda}) = \frac{2.24 e^{\frac{\overline{\lambda}^2}{2}}}{\sqrt{1 - \overline{\lambda}}} + \frac{e^{\overline{\lambda} + \overline{\lambda}^2}}{\sqrt{\pi}(1 - \overline{\lambda})}.$$

Noting that $c_x(\overline{\lambda})$ is increasing in $\overline{\lambda} \in [0, 1)$, we have, for all $0 \le x \le 0.606\sigma$,

$$\mathbf{P}(X_n > x\sigma) = \left(\Theta(x) + \theta \frac{c_x}{\sigma} \right) \inf_{\lambda \ge 0} \mathbf{E} e^{\lambda(X_n - x\sigma)},$$

where

$$c_x = \frac{2.24 e^{\frac{t^2}{2}}}{\sqrt{1 - t}} + \frac{e^{t + t^2}}{\sqrt{\pi}(1 - t)}$$

with t defined in (53). In particular, using the inequality $\mathbf{P}(X_n \ge x\sigma) \le \inf_{\lambda \ge 0} \mathbf{E} e^{\lambda(X_n - x\sigma)} \le H_n(x, \sigma)$ (cf. (35)), we obtain, for all $x \ge 0$,

$$\mathbf{P}(X_n > x\sigma) \le \left(\left(\Theta(x) + \frac{c_\alpha}{\sigma} \right) \wedge 1 \right) H_n(x, \sigma).$$

This completes the proof of Theorem 3. $\square$

References

1. Bahadur, R. and Rao, R., 1960. On deviations of the sample mean. *Ann. Math. Statist.*, **31**, 1015–1027.
2. Bennett, G., 1962. Probability inequalities for sum of independent random variables. *J. Amer. Statist. Asso.* **57**, No. 297, 33–45.
3. Bentkus, V., 2002. A remark on Bernstein, Prokhorov, Bennett, Hoeffding, and Talagrand inequalities. *Lithuanian Math. J.* **42**, No. 2, 262–269.
4. Bentkus, V., 2004. On Hoeffding's inequality, *Ann. Probab.* **32**, No. 2, 1650–1673.
5. Bercu, B. and Rouault, A., 2006. Sharp Large Deviations for the Ornstein–Uhlenbeck Process, *Theory Probab. Appl.*, **46**, No. 1, 1–19.
6. Bercu, B. and Touati, A., 2008. Exponential inequalities for self-normalized martingales with applications, *Ann. Appl. Probab.*, 1848–1869.
7. Bercu, B. et al., 2011. Sharp large deviations for the fractional Ornstein-Uhlenbeck process, *Theory Probab. Appl.*, **55**, No. 4, 575–610.
8. Cramér, H., 1938. Sur un nouveau théorème-limite de la théorie des probabilités. *Actualite's Sci. Indust.* **736**, 5–23.
9. Delyon, B., 2009. Exponential inequalities for sums of weakly dependent variables. *Electron. J. Probab.* **14**, 752–779.
10. Dembo, A., Zeitouni, O., 1998. *Large deviations techniques and applications* (2nd ed.). Springer, New York.
11. Deuschel, J.D., Stroock, D.W., 1989. *Large deviations.* Boston: Academic Press.
12. Doukhan, P., and Neumann, M. H., 2007. Probability and moment inequalities for sums of weakly dependent random variables, with applications, *Stochastic Process. Appl.*, **117** 878–903.
13. Eaton, M.L., 1974. A probability inequality for linear combination of bounded randon variables. *Ann. Statist.* **2**, No. 3, 609–614.
14. Fan, X., Grama, I. and Liu, Q., 2012. Hoeffding's inequality for supermartingales. *Stochastic Process. Appl.* **122**, 3545–3559.
15. Feller, W., 1971. *An introduction to probability theory and its applications.* J. Wiley and Sons.
16. Grama, I. and Haeusler, E., 2000. Large deviations for martingales via Cramer's method. *Stochastic Process. Appl.* **85**, 279–293.
17. Györfi, L., Harremöes, P. and Tusnády, G., 2012. Some refinements of large deviation tail probabilities. *arXiv:1205.1005v1 [math.ST].*
18. Haeusler, E., 1984. An exact rate of convergence in the functional central limit theorem for special martingale difference arrays. *Probab. Theory Relat. Fields* **65**, No. 4, 523–534.
19. Hahn, M. G. and Klass, M. J., 1997. Approximation of partial sums of arbitrary i.i.d. random varibales and the precision of the usual exponential upper bound. *Ann. Probab.* **25**, 1451–1470.
20. Hoeffding, W., 1963. Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.* **58**, 13–30.
21. Ledoux, M. and Talagrand, M., 1996. *Probability in Banach spaces.* Springer-Verlag.
22. Nagaev, S.V., 1979. Large deviations of sums of independent random variabels. *Ann. Probab.* **7**, No. 5, 745–789.
23. Nagaev, S.V., 2002. Lower bounds for the probabilities of large deviations of sums of independent random variables. *Theory Probab. Appl.* **46**, no 1, 79–102; no 4, 728–735.
24. Nagaev, S.V., 2003. On probability and moment inequalities for supermartingales and martingales. *Acta. Appl. Math.* **79**, 35–46.
25. Nagaev, S.V., 2007. On probability and moment inequalities for supermartingales and martingales. *Acta. Appl. Math.* **97**, 151–162.
26. Nagaev, S.V. and Chebotarev, V.I., 2011. On an estimate for the closeness of the binomial distribution to the the normal distribution. *Dokl. Math.* **83**, no 1, 19–21.
27. Rio, E., 2002. A Bennet type inequality for maxima of empirical processes. *Ann. Inst. H. Poincaré Probab. Statist.* **38** 1053–1057.
28. Rio, E., 2012. About the rate function in Talagrand's inequality for empirical processes. *C.R. Acad. Sci. Paris,* Ser. I. **350** 303–305.
29. Rozovky, L.V., 2003. A lower bound of large-deviation probabilities for the sample mean under the Cramér condition. *J. math. Sci.* **118**, No 6.
30. Rozovky, L.V., 2005. large deviation probabilities for some classes of distributions statisfying the Cramér condition. *J. math. Sci.* **128**, No 1.
31. Petrov, V.V., 1975. *Sums of Independent Random Variables.* Springer-Verlag. Berlin.
32. Petrov, V. V., 1995. *Limit Theorems of Probability Theory.*. Oxford University Press, Oxford.
33. Petrov, V.V. and Robinson, J., 2008. Large deviations for sums of independent non identical distributed random variables. *Comm. Statist. Theory Methods* **37**, 2984–2990.
34. Pinelis, I. and Utev, S. A., 1989. Exact exponential bounds for sums of independent random variables. *Theory Probab. Appl.* **34** 340–346.

35. Pinelis, I., 2006. Binomial uper bounds on generalized moments and tail probabilities of (super)martingales with differences bounded from above. *High Dimensional probab.* **51**, 33–52.
36. Shevtsova, I.G., 2010. An improvement of convergence rate estimates in the Lyapunov theorem. *Doklady. Math.* **82** 862–864.
37. Talagrand, M., 1995. The missing factor in Hoeffding's inequalities. *Ann. Inst. H. Poincaré Probab. Statist.* **31**, 689–702.
38. Talagrand, M., 1996b. A new look at independence. *Ann. Probab.* **22**, 1–34.
39. Talagrand, M., 1996a. New concentration inequalities in product spaces. *Invent. Math.* **126**, 505–563.
40. van de Geer, S., 2002. On Hoeffding's inequality for dependent random variables. In: Empirical Process Techniques for Dependent Data, eds. H. Dehling, T. Mikosch and M. Sorensen 161-170. Birkhauser, Boston.
41. van de Geer, S., 1995. Exponential inequalities for martingales, with application to maximum likelihood estimation for counting process. *Ann. Statist.*, **23**, 1779–1801.
42. van de Geer, S. and Lederer, J., 2011. The Bernstein-Orlicz norm and deviation inequalities. *Probab. Theory Relat. Fields*, DOI 10.1007/s00440-012-0455-y, 1–26.